

Takahiro Mizusaki

Senshu University, Japan

Pfaffian, Onishi formula and Wick's theorem

Since the pioneering work of Robledo [1], the Pfaffian has been extensively studied in nuclear physics. With the use of the Grassmann integrals, his work gives an excellent answer to the sign-problem in the overlap between HFB wave functions (Onishi formula [2] proposed in half a century ago). After Robledo's work, I have also studied some extensions [3] and applications. Here I will talk about the following topics:

1. Basics of Pfaffian and matrix elements of quantum many-body system
2. Pfaffian and sign-problem of Onishi formula
 - Pfaffian formula for overlap of HFB wave functions
 - Origin of the sign-problem and linked cluster theorem
3. Pfaffian and Wick's theorem

I will show some examples of Pfaffian in other fields of quantum many-body physics.

[1] L.M. Robledo, Phys. Rev. C 79 (2009), 021302(R).

[2] N. Onishi, S. Yoshida, Nucl. Phys. 80 (1966) 367.

[3] M. Oi, T. Mizusaki, Phys. Lett. B 707 (2012) 305, T. Mizusaki, M. Oi, Phys. Lett. B 715 (2012) 219, T. Mizusaki, M. Oi, F. Chen, Y. Sun, Phys. Lett. B 725 (2013) 175, T. Mizusaki, M. Oi, N. Shimizu, Phys. Lett. B 779 (2018) 237.

Mercredi 6 Jun. 2018, 11h30

IPN, Bât. 100, Salle A015