

SÉMINAIRE du GROUPE THÉORIE



INSTITUT DE PHYSIQUE NUCLÉAIRE

Groupe de Physique Théorique

Bât. 100, F-91406 ORSAY CEDEX

Tél (33)-(0)1-6915-7330 - Fax (33)-(0)1-6915-7748



Anna Zdeb

CEA

Cluster radioactivity of super-heavy nuclei.

An exotic type of decay - cluster emission - was observed in actinides region in 1980's [1, 2, 3]. In this process a light nucleus, but heavier than alpha particle is emitted, while the heavy mass residue is close to doubly magic lead in all observed cases. Microscopic description of this phenomenon treated as a very asymmetric fission has been successfully performed within HFB model with Gogny interactions D1S [4]. The fission valley on the potential energy surface has been found and fission fragments have been identified as cluster radioactivity products. Recently the characteristic very asymmetric fission valleys have been also found in some super-heavy nuclei. It has been shown that cluster radioactivity plays a nonnegligible role in this part of nuclear chart [5].

[1] H. J. Rose and G. A. Jones, *Nature (London)* 307, 245 (1984).

[2] A. Sandulescu, D. N. Poenaru, and W. Greiner, *Sov. J. Part. Nucl.* 11, 528 (1980).

[3] R. Bonetti and A. Guglielmetti, in *Heavy Elements and Related Phenomena*, Vol. II, edited by W. Greiner and R. K. Gupta (World Scientific, Singapore, 1999), p. 643.

[4] M. Warda, J.M. Robledo, *Phys. Rev. C* 84, 044608 (2011).

[5] M. Warda, A. Zdeb, L.M. Robledo, arXiv:1807.00342

Mercredi 16 Jan. 2019, 11h30

IPN, Bât. 100, Salle A015