Perspectives and Questions Meditations on the Future of Particle Physics Chris Quigg Fermilab & ENS

Séminaire commun, LPTENS/LPTHE · 02.04.2019

Supplemental reading: "Dream Machines," arXiv:1808.06036

CHF200 Note (2018)

... many scales



The importance of the 1-TeV scale

EW theory does not predict Higgs-boson mass Thought experiment: conditional upper bound

 W^+W^- , ZZ, HH, HZ satisfy *s*-wave unitarity, provided $M_H \lesssim (8\pi\sqrt{2}/3G_F)^{1/2} \approx 1 \text{ TeV}$

If bound is respected, perturbation theory is "everywhere" reliable If not, weak interactions among W^{\pm} , Z, H become strong on 1-TeV scale New phenomena (H or something else) are to be found around 1 TeV

Where is the next important scale?

(Higher energies needed to measure HHH, verify that H regulates $W_L W_L$)

 $\label{eq:Planck scale} \begin{array}{l} \text{Planck scale} \sim 1.2 \times 10^{19} \ \text{GeV} \ (3+1\text{-d spacetime}); \\ \sim 1.6 \times 10^{-35} \ \text{m} \\ \\ \text{Unification scale} \sim 10^{15-16} \ \text{GeV} \end{array}$

 $\Lambda_{QCD} \sim$ scale of confinement, chiral symmetry breaking

At what scale are charged-fermion masses set (Yukawa couplings)? At what scale are neutrino masses set? Will new physics appear at $1 \times, 10 \times, 100 \times, \ldots$ EW scale? Might new phenomena appear at macroscopic scales? The Great Lesson of Twentieth-Century Science

The human scale of space and time is not privileged for understanding Nature, and may even be disadvantaged.

Renormalization group · Effective field theories Resolution and extent in time and distance Diversity and scale diversity in experimental undertakings

The discovery that the human scale is not preferred is as important as the discoveries that the human location is not privileged (Copernicus) and that there is no preferred inertial frame (Einstein), and will prove to be as influential. How to progress?

Explore the regions of the unknown, the unanswered questions

Try to divine where the secrets are hidden

Seek out soft spots in our current understanding, especially where the stories we tell are $unprincipled \equiv$ not founded on sound principles

Supersymmetry: + R-parity + μ problem + tame FCNC + . . .

Big-Bang Cosmology: + inflation + dark matter + dark energy + ...

Particle content, even gauge groups, of the Standard Model

Chris Quigg

Guiding Principles Symmetry (via Noether's Theorems) & Hidden Symmetry Poincaré Invariance Relativistic Quantum Field Theory Unitarity, Causality ?? Renormalizability ?? Working hypotheses: Gauge Symmetry Pointlike constituents Minkowski spacetime (for most purposes)

Questions about fundamentals

- Is Lorentz invariance exact?
- Are nature's laws the same at all times and places (accessible to us)?
- What is the domain of validity of local field theory?
- Can causality be violated?
- Is CPT a good symmetry?
- Do quarks and leptons show signs of compositeness? Are they made of more elementary constituents?
- Are there supplemental spacetime dimensions?

On-mass-shell accelerators

Large Hadron Collider Complex at CERN Fermilab Main Injector J-PARC Main Ring BEPC II (IHEP-Beijing) VEPP-2000 (BINP-Novosibirsk)

SuperKEKB (first collisions, 25.03.2019) Intensity improvement projects for ν physics (Fermilab, J-PARC) [Facility for Antiproton and Ion Research (Darmstadt)]

HL-LHC, promising 3000 fb⁻¹ at $\sqrt{s} \rightarrow 14$ TeV

Virtual accelerators

Japan: ILC, e^+e^- collisions initially at $\sqrt{s} = 250$ GeV HE-LHC (energy doubler for the LEP/LHC tunnel), pp at $\sqrt{s} \approx 27$ TeV CLIC-380, e^+e^- collisions initially up to $\sqrt{s} = 380$ GeV LHeC. to collide a 60-GeV e beam with the LHC p beam Electron-Ion Collider, developed at Brookhaven and JLab CERN Future Circular Colliders: 100-km tunnel, ee, hh, eh studies China: CEPC (e^+e^- Higgs factory) in large tunnel \rightsquigarrow SppC (Muon Accelerator Program & Low EMmittance Muon Accelerator)

What LHC has taught us about the Higgs Boson

Evidence is developing as it would for a "standard-model" Higgs boson Unstable neutral particle with $M_H = 125.18 \pm 0.16$ GeV Decays to W^+W^- , ZZ implicate H as agent of EWSB Decay to $\gamma\gamma$ as expected (loop-level) Indirect constraint on Γ_H Dominant spin-parity $J^P = 0^+$ $Ht\bar{t}$ coupling from gg fusion, $t\bar{t}H$ production link to fermion mass origin

 $au^+ au^-$ and $bar{b}$ at expected rates

Only third-generation fermion couplings observed; $\mu^+\mu^-$ constrained

Search-and-discovery phase \rightsquigarrow painstaking forensic investigation

Questions about EWSB and the Higgs Sector

- Is H(125) the only member of its clan? Might there be others—charged or neutral—at higher or lower masses?
- Does H(125) fully account for electroweak symmetry breaking? Does it match standard-model branching fractions to gauge bosons? Are absolute couplings to W and Z as expected in the standard model?
- Are all production rates as expected? Any surprise sources of H(125)?
- What accounts for the immense range of fermion masses?
- Is the Higgs field the only source of fermion masses?
 Are fermion couplings proportional to fermion masses? $\mu^+\mu^-$ soon?
 How can we detect $H \rightarrow c\bar{c}$? e^+e^- ?? (basis of chemistry)
- What role does the Higgs field play in generating neutrino masses?

More questions about EWSB and the Higgs Sector

- Can we establish or exclude decays to new particles? Does H(125) act as a portal to hidden sectors? When can we measure Γ_H ?
- Can we detect flavor-violating decays ($au^{\pm}\mu^{\mp}$, ...)?
- Do loop-induced decays $(gg, \gamma\gamma, \gamma Z)$ occur at standard-model rates?
- What can we learn from rare decays $(J/\psi \gamma, \Upsilon \gamma, \dots)$?
- Does the EW vacuum seem stable, or suggest a new physics scale?
- Can we find signs of new strong dynamics or (partial) compositeness?
- Can we establish the HHH trilinear self-coupling?
- How well can we test the notion that H regulates Higgs-Goldstone scattering, i.e., tames the high-energy behavior of WW scattering?
- Is the electroweak phase transition first-order?

See Dawson, Englert, Plehn, arXiv:1808.01324

More new physics on the TeV scale and beyond?

Before LHC, much informed speculation—but no guarantees—about what might be found, beyond keys to EWSB.

Many eyes were on supersymmetry or Technicolor to enforce $M_W \ll$ unification scale or Planck scale.

"WIMP miracle" pointed to the TeV scale for a dark matter candidate.

Some imagined that neutrino mass might be set on the TeV scale.

No direct sign of physics beyond the standard model has come to light.

Might first hints may come from precision measurements?

Have we misconstrued naturalness and the hierarchy problem?

Did the existence of two once-and-done candidate solutions to the hierarchy problem (supersymmetry and technicolor) lead us to view the discipline of naturalness too simplistically?



Available online at www.sciencedirect.com SCIENCE DIRECT.

Nuclear Physics B (Proc. Suppl.) 140 (2005) 3-19



www.elsevierphysics.com

The Origins of Lattice Gauge Theory

K.G. Wilson

Smith Laboratory, Department of Physics, The Ohio State University, 174 W. 18th Ave., Columbus, OH 43210

Cf. M. Dine, "Naturalness under Stress" G. F. Giudice. "The Dawn of the Post-Naturalness Era"

Chris Quigg

Perspectives and Questions . . .

The final blunder was a claim that scalar elementary particles were unlikely to occur in elementary particle physics at currently measurable energies unless they were associated with some kind of broken symmetry [23]. The claim was that, otherwise, their masses were likely to be far higher than could be detected. The claim was that it would be unnatural for such particles to have masses small enough to be detectable soon. But this claim makes no sense when one becomes familiar with the history of physics. There have been a number of cases where numbers arose that were unexpectedly small or large. An early example was the very large distance to the nearest star as compared to the distance to the Sun, as needed by Copernicus, because otherwise the nearest stars would have exhibited measurable parallax as the Earth moved around the Sun. Within elementary particle physics, one has unexpectedly large ratios of masses, such as the large ratio of the muon mass to the electron mass. There is also the very small value of the weak coupling constant. In the time since my paper was written, another set of unexpectedly small masses was discovered: the neutrino masses. There is also the riddle of dark energy in cosmology, with its implication of possibly an extremely small value for the cosmological constant in Einstein's theory of general relativity.

Questions about new physics on the TeV scale and beyond

- Are there new forces of a novel kind?
- Scan we find evidence of a dark matter candidate?
- Why is empty space so nearly massless? What is the resolution to the vacuum energy problem?
- Will "missing energy" events signal the existence of spacetime dimensions beyond the familiar 3 + 1?
- ② Can we probe dark energy in laboratory experiments?
- Can we find clues to the origin of electroweak symmetry breaking? Is there a dynamical origin to the "Higgs potential?"
- What separates the electroweak scale from higher scales?

More questions about new physics on the TeV scale and beyond

• Might we find indirect evidence for a new family of strongly interacting particles, such as those that are present in SUSY, by seeing a change in the evolution of $1/\alpha_s(Q^2)$?



- How can we constrain—or provide evidence for—light dark-matter particles or other denizens of the dark in high-energy colliders or beam-dump experiments?
- Does the gluon have heavy partners, indicating that QCD is part of a structure richer than SU(3)_c?

Flavor: the problem of identity

What makes an electron an electron, a top quark a top quark, ...?
We do not have a clear view of how to approach the diverse character of the constituents of matter

CKM paradigm: extraordinarily reliable framework in hadron sector

BUT—many parameters: no clue what determines them, nor at what energy scale they are set

Even if Higgs mechanism explains *how* masses and mixing angles arise, we do not know *why* they have the values we observe

Physics beyond the standard model!

Flavor: the problem of identity (continued) Parameters of the Standard Model

- 3 Coupling parameters, α_{s} , α_{em} , $\sin^2 \theta_{W}$
- 2 Parameters of the Higgs potential
- 1 Vacuum phase (QCD)
- 6 Quark masses
- 3 Quark mixing angles
- 1 CP-violating phase
- 3 Charged-lepton masses
- 3 Neutrino masses
- 3 Leptonic mixing angles
- 1 Leptonic CP-violating phase (+ Majorana phases?)

26⁺ Arbitrary parameters

Will we see or diagnose a break in the SM? Contrast Landscape

Questions concerning the problem of identity

- Can we find evidence of right-handed charged-current interactions? Is nature built on a fundamentally asymmetrical plan, or are the right-handed weak interactions simply too feeble for us to have observed until now, reflecting an underlying symmetry hidden by spontaneous symmetry breaking?
- What is the relationship of left-handed and right-handed fermions?
- Are there additional electroweak gauge bosons, beyond W^{\pm} and Z?
- Are there additional kinds of matter?
- Is charged-current universality exact?What about lepton-flavor universality?

More questions concerning the problem of identity

- What do generations mean? Is there a family symmetry?
- Where are flavor-changing neutral currents? In the standard model, these are absent at tree level and highly suppressed by the Glashow-Iliopouolos-Maiani mechanism. They arise generically in proposals for physics beyond the standard model, and need to be controlled.

And yet we have made no sightings! Why not?

- Can we find evidence for charged-lepton flavor violation?
- Why are there three families of quarks and leptons? (Is it so?)
- Are there new species of quarks and leptons? exotic charges?

Neutrinos . . .

Neutrinos oscillate among the three known species, ν_e, ν_μ, ν_τ (discovered with neutrinos from natural sources)

Accelerator-based experiments NO ν A and T2K \rightsquigarrow DUNE and Hyper-Kamiokande + new short-baseline experiments

> Tritium β -decay experiment KATRIN experiments that rely on reactors (JUNO) or natural sources (IceCube and KM3Net)

Puzzling results: LSND-MiniBooNE, "Reactor anomaly"

Some outstanding questions about neutrino physics

- What is the order of levels of the mass eigenstates v₁, v₂, v₃? It is known that the v_e-rich v₁ is the lighter of the "solar pair," with the more massive v₂. Does the v_e-poor v₃ lie above or below the other two (normal or inverted mass ordering)?
- What is the absolute scale of neutrino masses? KATRIN vs. Cosmo?
- What is the flavor composition of ν_3 ? Is it richer in ν_{μ} or ν_{τ} ?
- Is CP violated in neutrino oscillations? To what degree?
- Are neutrinos Majorana particles? While this issue is primarily addressed by searches for neutrinoless double- β decay, collider searches for same-sign lepton pairs also speak to it.
- Do three light (left-handed) neutrinos suffice?
- What is the nature of right-handed neutrinos?

More outstanding questions about neutrino physics

- Are there light sterile neutrinos? If so, how could they arise?
- Do neutrinos have nonstandard interactions, beyond those mediated by W^{\pm} and Z?
- Below the cosmic neutrino background?
 Each species, now: 56 cm⁻³ $T_{\nu} \approx 2 \text{ K} \approx 1.7 \times 10^{-4} \text{ eV}$
- Are all the neutrinos stable?
- Do neutrinos contribute appreciably to the dark matter of the Universe?
- Bow is neutrino mass a sign of physics beyond the standard model?
- Will neutrinos give us insight into the matter excess in the Universe (through leptogenesis)?

Consider a neutrino factory

A Neutrino Factory based on a muon storage ring could provide a very strong second act for the coming generation of accelerator-based neutrino experiments.

Beyond its application to oscillation experiments as an intense source with known composition, an instrument that delivered $10^{20} \nu$ per year could be a highly valuable resource for on-campus experiments.

Neutrino interactions on thin targets, polarized targets, or active targets could complement the nucleon-structure programs carried out in electron scattering at Jefferson Lab and elsewhere.

Eventually: Multi-TeV muon collider in the LHC tunnel??

Don't forget the strong interactions!

Heroic progress in perturbative and lattice methods

QCD could be complete, up to M_{Planck} (modulo strong CP problem) ... but that doesn't prove it must be

Prepare for surprises, such as (Breakdown of factorization) Free quarks / unconfined color New kinds of colored matter Quark compositeness Larger color symmetry containing SU(3)_c

Questions pertaining to QCD

- Why is isospin a good symmetry? What does it mean?
- Are there new phenomena within QCD?
 Multiple production beyond diffraction + short-range order?
 Long-range correlations in y (or η)? Unusual event structures?
- Will the expected high density of few-GeV partons lead to thermalization in *pp* collisions? What will be other consequences?
- ${\color{black} \bullet}$ How will the 1-d $\infty {\color{black} -} momentum$ frame parton-model break down?
- How will correlations among partons in a proton manifest themselves?
- Can we distinguish spatial configurations of partons within protons?
- What is the importance of intrinsic heavy flavors?
- Hadron body plans beyond qqq and $q\bar{q}$? $QQ\bar{q}\bar{q}$, $qqqQ\bar{Q}$, ...
- Gan we prove that QCD confines color?

Motivations for unified theories

Neutrality of atoms, balance of electron and proton charges

Quarks and leptons are spin- $\frac{1}{2}$ particles that come in matched sets as required by anomaly cancellation for a renormalizable SU(2)_L \otimes U(1)_Y theory

 $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ gauge couplings tend to converge at high scales

Questions about unified theories

- What is the relationship of quarks to leptons?
- Should we regard lepton number as the "fourth color?"
- Which quark doublet is matched with which lepton doublet?
- Are there new gauge interactions that link quarks with leptons?
- What is the (grand) unifying symmetry?
- What determines the low-energy gauge symmetries?
- What are the steps to unification? One more, or multiple?
- Is perturbation theory a reliable guide to coupling-strength unification?
- What sets the mass scale for the additional gauge bosons in a unified theory? ... for the additional Higgs bosons?

More questions about unified theories

- Is the proton unstable? How does it decay?
- Is neutron-antineutron oscillation observable?
- Can we detect the magnetic monopoles of unified theories?
- Are there millicharged particles?
 Other signs of additional U(1) gauge symmetries?
- e How can we incorporate gravity?
- Why is gravity so weak?
- To what scale does the inverse-square law of gravitation hold?
- What is the nature of spacetime?
 Is it emergent?
 How many dimensions?

A word about the astro/cosmo connection

We do not know what the Universe at large is made of We do not know the complete thermal history of the universe

We have not accounted for the predominance of matter over antimatter in the observed universe

We do not know what provoked inflation (if it happened) We do not know why the expansion of the universe is accelerating i.e., the origin of dark energy or a cosmological constant

Detection of gravitational radiation enriches multimessenger astronomy

Questions about the universe at large

- To what degree does the cosmological principle hold (homogeneous and isotropic universe)?
- Have we entered a new inflationary epoch?
- Is there a dynamical interplay between cosmological evolution and scalar-field "relaxion" dynamics (including H)?
- What is the "dark energy" equation of state? How will dark energy evolve with time (∝ a^{-(1+w)})? Dynamics or Λ? If Λ, what sets the scale?
- Are there any alternatives/complements to collisionless dark matter?
- How can technologies developed for accelerators advance the search for axions? How can we observe axions, dark photons, ...?

G. Bertone & T. M. P. Tait, "A new era in the search for dark matter," Nature 562, 51-56 (2018)

Chris Quigg

Tabletop precision experiments

Electric dipole moment d_e : CP/T violation $|d_e| < 1.1 imes 10^{-29} \ e \ {
m cm}$ ACME Collaboration. ThO $|d_e| < 1.3 imes 10^{-28}~e$ cm NIST. trapped ¹⁸⁰Hf ¹⁹F⁺ (SM phases: $d_e < 10^{-38} e$ cm)

(How) can we observe electric dipole moments of e, μ, p ?

"Tabletop" precision experiments

(Anti)proton magnetic moments: CPT test

$$\mu_{ar{
ho}} = -2.792\,847\,344\,1(42)\;\mu_{N}$$

VS.

$\mu_{P} = +2.792\,847\,344\,62(82)\,\mu_{N}$

BASE Collaboration @CERN Antiproton Decelerator

Exercise 1.

How should we respond if:

(a) The DAMA "seasonal variation" cannot be explained away? cf. COSINE-100

- (b) The LHC Higgs signal strength settles at $\mu = 1.17 \pm 0.03$? Or if Ht \overline{t} remains high?
- (c) The LHCb flavor anomalies persist?
- (d) The $(g-2)_{\mu}$ anomaly strengthens?
- (e) WIMP dark matter searches reach the neutrino floor?

... (extra credit)

Exercise 2.

Sketch five "small-scale" (you define) experiments with the potential to change our thinking about particle physics or related fields.

Exercise 3.

How would *you* assess the scientific potential (in view of cost and schedule) of

(a) The High-Luminosity LHC? (b) The High-Energy LHC? (c) A 100-TeV pp Collider (FCC-hh)? (d) A 250-GeV ILC? (e) A circular Higgs factory (FCC-ee or CEPC)? (f) A 380-GeV CLIC? (g) LHeC / FCC-eh? (or an electron-ion collider?) (h) A muon-storage-ring neutrino factory? (i) A multi-TeV muon collider? The instrument of your dreams?

Chris Quigg

Big dreams

Fermi's dream accelerator (1954) $E_{ m beam} = 5\,000$ TeV, $\$1.7 imes 10^{11}$



A THOUSAND TOV IN THE CENTER OF MASS:

INTRODUCTION TO HIGH ENERGY STORAGE RINGS⁴

J.D. Bjorken

Fermi National Accelerator Laboratory P.O. Box 500 Batavia, Illinois 60510

I. INTRODUCTION

These lectures must begin with an apology. Normally at schools such as this, one expects the lecturer to be an acknowledged expert on the subject matter he is discussing. Here this is not the case. Design of high energy proton storage rings is not exactly my <u>forte.</u> Why am I doing this? There are several reasons, short of mental lilness.

1. I want to learn this subject myself and there is no better way than trying to teach it. And Ferbel didn't stop me.

2. There needs to be a broader knowledge of accelerator physics in the elementary-particle community. Experimentalists at the storage rings find themselves especially closely coupled to their machine and its operation. And theorists can find interesting and challenging questions which lie at the frontier of the very active field of nonlinear mechanics.

3. Straightforward extrapolation of existing acceleration techniques would seem to lead to very large, expensive machines. While we may envision one, perhaps two generations of future accelerators using essentially existing techniques, the question of how to go beyond that is a difficult one. There seems to be a growing feeling that it is not too soon to start to face up to the problem. A look at the alternative--as we do here--can only provide stimulation.

Questions inspired by Big Dreams

- Suppose we could reach gradients of many GeV—even 1 TeV—per meter. How would we first apply that bit of magic, and what characteristics other than gradient would be required?
- If we could shrink multi-TeV accelerators, how might we shrink detectors that depend on particle interactions with matter?
- What could we do with a low-emittance, high-intensity muon source?
- What inventions would it take to accelerate beams of particles with picosecond lifetimes?
- Beams? How might we apply high-transmissivity crystal channeling?
- How would optimizations change if we could shape superconducting magnet coils out of biplanar graphene or an analogous material?

Two final questions (for now)!

What deep questions have been with us for so long that they are less prominent in "top-ten" lists than they deserve to be?

How are we prisoners of conventional thinking? How can we break out?