

Gianluca Stellin

IJCLab, Université Paris-Saclay, 91405 Orsay, France

Spectroscopy of even-even open-shell nuclei via self-consistent Gorkov-Greens function calculations

The fundamentals of the *ab-initio* Self-Consistent Gorkov Greens function (SCGGF) [1, 2] approach for the investigation of low-lying energy spectrum of the semi-magic even-even nuclei are presented. In the last decade, the SCGGF method has brought a significant renewal in the realm of *ab-initio* approaches to nuclear structure, marking a step forward in the knowledge of bulk nuclear properties of even-even nuclei, such as the ones lying along the Ar-Cr [3, 4] isotopic chains. The access to the one-particle propagator has allowed the study of ground and excited states of neighbouring odd-A isotopes [5, 6, 7]. Nonetheless, the prediction of excited energy levels and reduced electric and magnetic multipole transition probabilities calls for the introduction of the polarization propagator, previously not embedded in the $U(1)_Z \times U(1)_N$ symmetry breaking formalism. In quantum chemistry, present-day approaches for the description of the spectrum of medium-sized organic molecules [8, 9] are based on diagrammatic many-body Greens function theory applied to the polarization propagator at third order in the *algebraic diagrammatic construction* (ADC) approach [10, 11, 12, 13]. Another return of this study will be provided by the prediction of new shell closures in neutron-rich even-even nuclei, identified through the local maxima in the energy of the 2_1^+ state and in the related electric quadrupole transition probability, $B(E2, 0_1^+ \rightarrow 2_1^+)$ [14].

References

- [1] L.P. Gorkov, *Sov. Phys. JETP* **34**, 3, 505-508 (1958).
- [2] V. Somà, T. Duguet and C. Barbieri, *Phys. Rev. C* **84**, 064317 (2011).
- [3] V. Somà, A. Cipollone, C. Barbieri, P. Navrátil and T. Duguet, *Phys. Rev. C* **89**, 061301(R) (2014).
- [4] V. Somà, C. Barbieri, T. Duguet and P. Navrátil, *Eur. Phys. J. A* **57**, 135 (2021).
- [5] M. Rosenbusch et al., *Phys. Rev. Lett.* **114**, 202501 (2015).
- [6] S. Chen et al., *Phys. Rev. Lett.* **123**, 142501 (2019).
- [7] Y.L. Sun et al., *Phys. Lett. B* **802**, 135215 (2020).
- [8] P.H.P. Harbach, M. Wormit and A. Dreuw, *J. Chem. Phys.* **141**, 064113 (2014).
- [9] A. Dreuw and M. Wormit, *Comput. Mol. Sci.* **5**, 82-95 (2015).
- [10] J. Schirmer, *Phys. Rev. A* **26**, 2395-2416 (1982).
- [11] A.B. Trofimov, G. Stelter and J. Schirmer, *J. Chem. Phys.* **111**, 9982-9999 (1999).
- [12] J. Brand and L.S. Cederbaum, *Adv. Quantum Chem.* **38**, 65-120 (2000).
- [13] A.B. Trofimov, G. Stelter and J. Schirmer, *J. Chem. Phys.* **117**, 6402-6409 (2002).
- [14] I. Bentley, Y. Colón Rodríguez, S. Cunningham and A. Aprahamian, *Phys. Rev. C* **93**, 044337 (2016).

Thursday 20th March 2025, 14h00
IJCLab, Build. 100, Room A201