

Séminaire de Physique Nucléaire Théorique

Spectroscopy of even-even open-shell nuclei via self-consistent Gorkov-Green's function calculations

Gianluca Stellan

(IJCLab)

The fundamentals of the *ab-initio* Self-Consistent Gorkov Green's function (SCGGF) [1-2] approach for the investigation of low-lying energy spectrum of the semi-magic even-even nuclei are presented. In the last decade, the SCGGF method has brought a significant renewal in the realm of ab-initio approaches to nuclear structure, marking a step forward in the knowledge of bulk nuclear properties of even-even nuclei, such as the ones lying along the Ar-Cr [3-4] isotopic chains. The access to the one-particle propagator has allowed the study of ground and excited states of neighbouring odd-A isotopes [5-6]. Nonetheless, the prediction of excited energy levels and reduced electric and magnetic multipole transition probabilities calls for the introduction of the polarization propagator, previously not embedded in the $U(1)_Z \times U(1)_N$ symmetry breaking formalism. In quantum chemistry, present-day approaches for the description of the spectrum of medium-sized organic molecules [7-8] are based on diagrammatic many-body Green's function theory applied to the polarization propagator at third order in the *algebraic diagrammatic construction* (ADC) approach [9-13]. Another return of this study will be provided by the prediction of new shell closures in neutron-rich even-even nuclei, identified through the local maxima in the energy of the 2_1^+ state and in the related electric quadrupole transition probability, $B(E2, 0_1^+ \rightarrow 2_1^+)$ [14].

[1] L.P. Gorkov, Sov. Phys. JETP 34, 3, 505-508 (1958). [2] V. Somà, T. Duguet and C. Barbieri, Phys. Rev. C 84, 064317 (2011). [3] V. Somà, A. Cipollone, C. Barbieri, P. Navrátil and T. Duguet, Phys. Rev. C 89, 061301(R) (2014). [4] V. Somà, C. Barbieri, T. Duguet and P. Navrátil, Eur. Phys. J. A 57, 135 (2021). [5] M. Rosenbusch et al., Phys. Rev. Lett. 114, 202501 (2015). [6] S. Chen et al., Phys. Rev. Lett. 123, 142501 (2019). [7] Y.L. Sun et al., Phys. Lett. B 802, 135215 (2020). [8] P.H.P. Harbach, M. Wormit and A. Dreuw, J. Chem. Phys. 141, 064113 (2014). [9] A. Dreuw and M. Wormit, Comput. Mol. Sci. 5, 82-95 (2015). [10] J. Schirmer, Phys. Rev. A 26, 2395-2416 (1982). [11] A.B. Trofimov, G. Stelter and J. Schirmer, J. Chem. Phys. 111, 9982-9999 (1999). [12] J. Brand and L.S. Cederbaum, Adv. Quantum Chem. 38, 65-120 (2000). [13] A.B. Trofimov, G. Stelter and J. Schirmer, J. Chem. Phys. 117, 6402-6409 (2002). [14] I. Bentley, Y. Colón

Rodríguez, S. Cunningham and A. Aprahamian, Phys. Rev. C 93, 044337 (2016).

Jeudi 20 Mars 2025

14 :00

IJCLab, Bât. 100, Salle Room A201