

Séminaire de Physique Nucléaire Théorique

Advances in microscopic modeling of $(n, xn\gamma)$ reactions for actinides

Marc Dupuis

(CEA, DAM, DIF)

Recent developments in nuclear structure approaches offer a great mean to improve various aspects of nuclear reaction modeling and to further understand reaction mechanisms from a microscopic point of view. Recently, direct and pre-compound nucleon emission, for nucleon induced reaction on spherical and axially deformed nuclei, have been successfully modeled [1] using a description of target states provided by fully consistent axially-symmetric deformed quasi-particle random-phase approximation (QRPA) calculations [2]. Direct inelastic scattering to target excitations built from one-phonon QRPA states accounted simultaneously for direct inelastic scattering to discrete states, and pre-equilibrium emission as far as second order processes, that involve more complex excitations such as two-phonon states, and multiple emission remain negligible. The QRPA nuclear structure approach has also been applied recently to determine, for a large pannel of even-even nuclei, E1 and M1 photon strength functions [3], that play a key-role in the modeling of statistical reactions.

We will review the status on the ongoing work on direct/pre-compound neutron emission for neutron induced reaction below 20 MeV for even-even actinides. Target states are described as rotational bands built from each state in the target intrinsic frame, described as QRPA one-phonon excitation of the intrinsic correlated ground state. QRPA excitations which display a collective character can thus be viewed as vibrational band heads. Couplings between states of the GS band and states belonging to an excited band are accounted for within a coupled channel framework.

Our approach is then applied to the modeling of $(n, n'\gamma)$ reactions and for both intra- and inter-band gamma transitions [4]. For these reactions, the role played by the present microscopic approach for direct/pre-equilibrium emission is discussed. We finally focus on the impact on the determination of $(n, n'\gamma)$ cross sections of newly calculated QRPA E1- and M1-photon strength functions, that enter the description of statistical decay from compound nucleus states in the continuum.

[1] M. Dupuis, E. Bauge, S. Hilaire, F. Lechaftois, S. Péru, N. Pillet and C. Robin, Eur. Phys. J. A, 51 12 (2015) 168.

- [2] S. P  ru, G. Gosselin, M. Martini, M. Dupuis, S. Hilaire, and J.-C. Devaux, Phys. Rev. C 014314 (2011).
- [3] S. Goriely, S. Hilaire, S. P  ru, M. Martini, I. Deloncle, and F. Lechaftois Phys. Rev. C 94, 044306 (2016) ; M. Martini, S. P  ru, S. Hilaire, S. Goriely, and F. Lechaftois, Phys. Rev. C 94, 014304 (2016).
- [4] M. Dupuis, S. Hilaire, S. P  ru, E. Bauge, M. Kerveno, P. Dessagne and G. Henning EPJ Web Conf., 146 (2017) 12002.

Jeudi 26 Juillet 2018

11 :30

IJCLab, B  t. 100, Salle B  timent 100, Salle A015