

Université Paris-Saclay  
IJCLab  
(Laboratoire de Physique des 2 Infinis Irène Joliot-Curie)  
Bât. 100, F-91405 Orsay

## Séminaire de Physique Nucléaire Théorique

### Insights into the reactions and structure of light nuclei from Effective Field Theory.

D. R. Phillips

*(Ohio University*

*Technische Universität Darmstadt)*

In this talk I will show how an Effective Field Theory for weakly-bound few-body systems ("Halo EFT") aids the modeling and understanding of light nuclei in the  $p$ -shell. After introducing Halo EFT I will discuss two examples of its use. First, I will describe our recent analysis of the reaction  ${}^3\text{He}(\alpha, \gamma)$  [1]. In this case we take  ${}^3\text{He}$  and  ${}^4\text{He}$  as the effective degrees of freedom and work to next-to-leading order in the EFT. We use a Bayesian analysis to perform the extrapolation of higher-energy data to solar energies and find a markedly smaller uncertainty to previous evaluations, as well as significant constraints on  ${}^3\text{He}$ - ${}^4\text{He}$  scattering parameters. Second, I will demonstrate that a three-body ( $\alpha$ -neutron-proton) model of  ${}^6\text{Li}$  exhibits a correlation between the deuteron- $\alpha$  scattering length and the  ${}^6\text{Li}$  binding energy [2]. Such correlations appear generically in three-body systems with weak binding (cf. the Phillips line in the  $A = 3$  system) and can be understood as a consequence of the requirement that the EFT be renormalized. The emergence of such a correlation in the presence of  $p$ -wave nucleon- $\alpha$  interactions is particularly striking, as is the fact that it is rather insensitive to the actual values of the  $p$ -wave phase shifts. I will close by discussing the implications of this finding for model and EFT treatments of  ${}^6\text{He}$  and  ${}^6\text{Li}$ .

**Mercredi 29 Mai 2019**

**11 :30**

**IJCLab, Bât. 100, Salle Bâtiment 100, Salle A015**